

ON A SUBPOSET OF THE TAMARI LATTICE

SEBASTIAN A. CSAR, RIK SENGUPTA, AND WARUT SUKSOMPONG


ABSTRACT. We explore some of the properties of a subposet of the Tamari lattice introduced by Pallo, which we call the comb poset. We show that three binary functions that are not well-behaved in the Tamari lattice are remarkably well-behaved within an interval of the comb poset: rotation distance, meets and joins, and the common parse words function for a pair of trees. We relate this poset to a partial order on the symmetric group studied by Edelman.

1. INTRODUCTION

The set $\mathbb{T}_n$ of all full binary trees with n leaves, or parenthesizations of n letters, has been well-studied, and carries much structure. Its cardinality $|\mathbb{T}_n|$ is the $(n-1)^{\text{th}}$ Catalan number

$$C_{n-1} = \frac{1}{n} \binom{2n-2}{n-1}.$$


The *rotation graph*, $\mathcal{R}_n$, is the graph with vertex set $\mathbb{T}_n$, in which edges correspond to a local change in the tree called a *rotation*, corresponding to changing a single parenthesis pair in the parenthesization. This graph $\mathcal{R}_n$ forms the vertices and edges of an $(n-2)$ -dimensional convex polytope called the *associahedron*, K_{n+1} . If we direct the edges of $\mathcal{R}_n$ in a certain fashion, we obtain the Hasse diagram for the well-studied *Tamari lattice*, $\mathcal{T}_n$, on $\mathbb{T}_n$, shown below for $n = 4$.



The Tamari lattice has many properties, but it has certain deficiencies. For instance, it is not ranked. Although one can encode the Tamari order by componentwise comparison of the *bracketing vectors* $\langle T \rangle \in \{0, 1, \dots, n-2\}^{n-1}$ of $T \in \mathcal{T}_n$, introduced by Huang and Tamari in [3], only the meet is given by the componentwise minimum of these bracketing vectors; the join cannot be characterized similarly. Furthermore, computing the *rotation distance* $d_{\mathcal{R}_n}(T_1, T_2)$ between two trees T_1, T_2 in the graph $\mathcal{R}_n$ does not appear to follow easily from knowing their meet and join in the Tamari lattice.

There is another binary function on $\mathbb{T}_n$ that becomes relevant in the light of an approach to the Four Color Theorem suggested by Kauffman in [4] and explored more recently by Cooper, Rowland and Zeilberger in [1]: the size of the set $\text{ParseWords}(T_1, T_2)$ consisting of all words $w \in \{0, 1, 2\}^n$ which are *parsed* by both T_1 and T_2 . Here, a word w is parsed by T if the labeling of the leaves of T by $w_1, w_2, \dots, w_n$ from left to right extends to a proper 3-coloring with colors $\{0, 1, 2\}$ of *all* $2n - 1$ vertices in T , such that no two children of the same vertex have the same label and such that no parent and child share the same label. Kauffman showed that the Four Color Theorem is equivalent to the statement that for all n and all $T_1, T_2 \in \mathbb{T}_n$, one has $|\text{ParseWords}(T_1, T_2)| \geq 1$.

This last application to the Four Color theorem motivated us to investigate a poset $\mathcal{C}_n$ on the set $\mathbb{T}_n$, which we call the *(right) comb order*, a weakening of the Tamari order. Pallo first defined $\mathcal{C}_n$ in [7], where he proved that it is a meet-semilattice having the same bottom element as $\mathcal{T}_n$, called the right comb tree and denoted $\text{RCT}(n)$. The solid edges in the diagram below form the Hasse diagram of $\mathcal{C}_4$. The dashed edge lies in $\mathcal{T}_4$ but not in $\mathcal{C}_4$.



While the comb order $\mathcal{C}_n$ is a meet-semilattice whose meet $\wedge_{\mathcal{C}_n}$ does not in general coincide with the Tamari meet $\wedge_{\mathcal{T}_n}$, it fixes several deficiencies of $\mathcal{T}_n$ noted above:

- $\mathcal{C}_n$ is ranked, with exactly $\binom{n+r-2}{r} - \binom{n+r-2}{r-1}$ elements of rank r (see Theorem 3.2).
- $\mathcal{C}_n$ is *locally distributive*; each interval forms a distributive lattice (see Corollary 2.11(i)).
- If T_1 and T_2 have an upper bound in $\mathcal{C}_n$ (or equivalently, if they both lie in some interval), the meet $T_1 \wedge_{\mathcal{C}_n} T_2$ and join $T_1 \vee_{\mathcal{C}_n} T_2$ are easily described, either in terms of intersection or union of reduced parenthesizations (see Corollary 2.11(i)), or by componentwise minimum or maximum of their bracketing vectors (see Theorem 5.4). These operations also coincide with the Tamari meet $\wedge_{\mathcal{T}_n}$ and Tamari join $\vee_{\mathcal{T}_n}$ (see Corollary 5.5).
- When trees T_1, T_2 have an upper bound in $\mathcal{C}_n$, one has (see Theorem 4.3)

$$\begin{aligned} d_{\mathcal{C}_n}(T_1, T_2) &= \text{rank}(T_1) + \text{rank}(T_2) - 2 \cdot \text{rank}(T_1 \wedge_{\mathcal{C}_n} T_2) \\ &= 2 \cdot \text{rank}(T_1 \vee_{\mathcal{C}_n} T_2) - (\text{rank}(T_1) + \text{rank}(T_2)) \\ &= \text{rank}(T_1 \vee_{\mathcal{C}_n} T_2) - \text{rank}(T_1 \wedge_{\mathcal{C}_n} T_2), \end{aligned}$$

where, for any $T \in \mathbb{T}_n$, $\text{rank}(T)$ refers to the rank of T in $\mathcal{C}_n$.

- Furthermore, for T_1, T_2 having an upper bound in $\mathcal{C}_n$, one has (see Theorem 7.9)

$$\text{ParseWords}(T_1, T_2) = \text{ParseWords}(T_1 \wedge_{\mathcal{C}_n} T_2, T_1 \vee_{\mathcal{C}_n} T_2),$$

with cardinality $3 \cdot 2^{n-1-k}$, where $k = \text{rank}(T_1 \vee_{\mathcal{C}_n} T_2) - \text{rank}(T_1 \wedge_{\mathcal{C}_n} T_2)$ (see Theorem 7.7).

Lastly, Section 6 discusses a well-known order-preserving surjection from the (right) weak order on the symmetric group $\mathfrak{S}_n$ to the Tamari poset $\mathcal{T}_{n+1}$ and its restriction to an order-preserving surjection from $\mathcal{E}_n$ to $\mathcal{C}_{n+1}$ (where $\mathcal{E}_n$ is a subposet of the weak order considered by Edelman in [2]). Furthermore, this surjection is a distributive lattice morphism on each interval of $\mathcal{C}_{n+1}$ (see Theorem 6.9).

Because we will be mainly confining our attention for the rest of this paper to the poset $\mathcal{C}_n$, we will drop the subscripts from $\wedge$, $\vee$, $>$ and $<$ when we mean meet, join, greater than, and less than in $\mathcal{C}_n$ respectively. Furthermore, we will use $\text{rank}(T)$ to denote the rank of T in $\mathcal{C}_n$. Much of our notation in Section 7 is from [1].

2. THE COMB POSET AND DISTRIBUTIVITY

Definition 2.1. For each binary tree $T \in \mathbb{T}_n$, consider the usual parenthesization of its leaves $a_1, a_2, \dots, a_n$. Then, delete all pairs of parentheses that enclose the leaf a_n . Call the resulting parenthesization the *reduced parenthesization* of T , denoted RP_T . Define an *element* of RP_T to be either an unparenthesized leaf in RP_T , or any pair of parentheses J in RP_T (together with all enclosed leaves and internal parenthesizations) which is *not* enclosed by some other pair of parentheses in RP_T .

Example. The reduced parenthesization of the following tree is $a_1((a_2a_3)a_4)(a_5a_6)a_7$. The reduced




FIGURE 1.

parenthesization of this tree has four elements, given by a_1 , $((a_2a_3)a_4)$, (a_5a_6) , and a_7 . Note that a_n by itself is always an element of the reduced parenthesization of any tree in $\mathbb{T}_n$.


Proposition 2.2. *Given the leaves $a_1, a_2, \dots, a_n$ in that order, a parenthesization of $a_1, a_2, \dots, a_n$ is the reduced parenthesization for some tree $T \in \mathbb{T}_n$ iff the two following conditions hold: a_n is not enclosed by any parenthesis pair, and each pair of parentheses encloses precisely two factors within it (which we shall call the left factor and the right factor).*

Remark. All n -leaf binary trees have a unique reduced parenthesization, since there is a bijection between the full parenthesization of a tree T and its reduced parenthesization. The full parenthesization is recovered by pairing the two rightmost elements of RP_T successively.

Remark. If two pairs of parentheses J_1 and J_2 enclose the same two factors in the same order, then the internal parenthesization of J_1 and J_2 must be the same. Definition 2.1 then implies that, if $RP_{T_1} \subset RP_{T_2}$, then each of the parenthesis pairs common to the two reduced parenthesizations also has the *same* factors in both RP_{T_1} and RP_{T_2} .

Definition 2.3. For $n \geq 2$, the *right comb tree* of order n , denoted by $\text{RCT}(n) \in \mathbb{T}_n$, is the n -leaf binary tree corresponding to the “empty” reduced parenthesization $a_1a_2 \cdots a_n$. Similarly, the *left comb tree* of order n is defined as the n -leaf binary tree corresponding to the reduced parenthesization $((\cdots((a_1a_2)a_3) \cdots)a_{n-2})a_{n-1})a_n$.

Example. $\text{RCT}(5)$, the right comb tree of order 5, is shown below. The nodes labeled $a_1, \dots, a_5$ are the leaves of the tree, and $b_6, \dots, b_9$ are the internal vertices. Note that the structure of the left comb tree of order 5 is given by the reflection of the right comb tree about the vertical axis.



Definition 2.4. Define the (*right*) *comb poset* of order n to be the poset whose elements are given by elements from $\mathbb{T}_n$, with $T_1 \leq T_2$ iff $RP_{T_1} \subseteq RP_{T_2}$, i.e. each parenthesis pair in RP_{T_1} appears in RP_{T_2} . Denote the right comb poset of order n by $\mathcal{C}_n$.

Remark. Since the $RCT(n)$ corresponds to the “empty” parenthesization $a_1 a_2 \cdots a_n$, it is the unique minimal element of $\mathcal{C}_n$.

Example. The Hasse diagram of the right comb poset of order 5 is shown in Figure 2. For the sake of a cleaner diagram, the leaf a_i is labeled by i in Figure 2 for $i \in \{1, 2, 3, 4, 5\}$.




FIGURE 2. The Hasse diagram of $\mathcal{C}_5$

Proposition 2.5. *There is an order-preserving involution on $\mathcal{C}_n$.*

Proof. For any tree T , take RP_T , and construct a new parenthesization $RP_{T'}$ as follows. For every parenthesis pair in RP_T which encloses leaves a_i through a_j , take $RP_{T'}$ to have a parenthesis pair enclosing leaves a_{n-j} through a_{n-i} . It is not hard to see (using Proposition 2.2) that $RP_{T'}$ corresponds to a tree T' . Define π to be the map that takes T to T' as described above. Then, π is an order preserving involution on $\mathcal{C}_n$. $\square$

Definition 2.6. For $n \geq 2$, the *right arm* of a tree $T \in \mathbb{T}_n$ is the connected acyclic graph induced by the vertices of T that lie in the left subtree of no other vertex in T .

Definition 2.7. Given an n -leaf binary tree, T , one obtains its *pruned tree* by deleting all the leaves from T , obtaining an “incomplete” tree on $n - 1$ vertices.




FIGURE 3.

It is well known that the operation of “pruning” is a bijection between n -leaf binary trees and (possibly incomplete) binary trees with $n - 1$ vertices.

Definition 2.8. For a tree $T \in \mathbb{T}_n$, the *reduced pruned poset* of T , denoted P_T , is the poset of pairs of parentheses in RP_T ordered by inclusion. Its Hasse diagram is obtained by pruning T , removing the right arm and removing edges incident to the right arm.

Example. Consider the tree of Figure 1, given by reduced parenthesization $a_1((a_2a_3)a_4)(a_5a_6)a_7$. Figure 3 depicts its “pruned” form and the corresponding reduced pruned poset P_T .

Proposition 2.9. For any tree $T \in \mathbb{T}_n$, the elements other than leaf a_n in RP_T correspond precisely to the dangling left subtrees of the vertices on the right arm of T , taken in order, with the leftmost element corresponding to the left subtree of the root.

Proposition 2.10. For any $T \in \mathbb{T}_n$, the interval $[\text{RCT}(n), T]_{\mathcal{C}_n}$ is isomorphic to the lattice of order ideals in the reduced pruned poset of T , ordered by inclusion. In other words, for any tree T ,

$$[\text{RCT}(n), T]_{\mathcal{C}_n} \cong J(P_T)$$

Proof. One has a natural map $J(P_T) \rightarrow [\text{RCT}(n), T]$, given by $I \mapsto S$, where S is the tree with RP_S having precisely the parentheses in I . The definition of the order on $\mathcal{C}_n$ ensures this map is both well-defined and order-preserving. Furthermore, this map has an inverse $[\text{RCT}(n), T] \rightarrow J(P_T)$ given by $S \mapsto \{\text{parentheses of } RP_S\}$, which is again order-preserving. $\square$

This proposition yields a number of immediate corollaries.

Corollary 2.11.

- (i) Any interval in $\mathcal{C}_n$ is a distributive lattice, with the reduced parenthesizations of the join and meet of trees T_1 and T_2 in an interval given by the ordinary union and intersection of parenthesis pairs from RP_{T_1} and RP_{T_2} .
- (ii) In $\mathcal{C}_n$, T_1 covers T_2 iff RP_{T_1} can be obtained from RP_{T_2} by adding one parenthesis pair.
- (iii) $\mathcal{C}_n$ is a ranked poset, with the rank of any tree T in $\mathcal{C}_n$ given by the number of parenthesis pairs in RP_T .
- (iv) For any two trees T_1 and T_2 that are in the same interval of $\mathcal{C}_n$, we have
$$\text{rank}(T_1) + \text{rank}(T_2) = \text{rank}(T_1 \wedge T_2) + \text{rank}(T_1 \vee T_2)$$
- (v) For any tree $T \in \mathbb{T}_n$ of rank k , the length of the right arm of T is $n - 1 - k$.

Note. In the remainder of this paper, we shall consider only the right comb poset of order n . Analogous results hold for left comb poset by symmetry.

Remark. The covering relation described in Corollary 2.11(ii) corresponds to a left rotation centered along the right arm of a tree, so-called right arm rotation, precisely the covering relation used by Pallo in [7] to define the poset $(B_n, \rightsquigarrow^*)$, which he showed to be a meet-semilattice [7, Lemma 3].

3. RANK SIZES IN THE COMB POSET

In this section, we will prove some enumerative properties of the ranks of $\mathcal{C}_n$. To simplify notation, let Q_i denote the i^{th} rank of $\mathcal{C}_n$.

Proposition 3.1. *For $0 \leq i \leq n - 2$, every tree in Q_i is covered by precisely $n - 2 - i$ trees.*

Proof. This fact follows from the definition of rotation, and the observation that a tree in rank Q_i has, by Corollary 2.11(v), a right arm of length $n - 1 - i$, i.e. $n - i$ vertices. $\square$

Theorem 3.2. *For $n \geq 3$, $\mathcal{C}_n$ is a ranked poset. A tree is a maximal element of $\mathcal{C}_n$ iff it is of rank $n - 2$ in $\mathcal{C}_n$ (or equivalently, from Corollary 2.11(v), iff its right arm has length 1). In particular, the left comb tree is in the maximal rank of $\mathcal{C}_n$. Furthermore, for $0 \leq r \leq n - 2$, the number of elements in rank r of $\mathcal{C}_n$ is*

$$|Q_r| = \binom{n+r-2}{r} - \binom{n+r-2}{r-1}.$$

Before the proof of the theorem, recall Lagrange Inversion. The proof may be found in [10, Theorem 5.1.1], among other places. Denote the coefficient of x^n in the power series function $\psi(y)$ by $[x^n]\psi(y)$.

Lemma 3.3 (Lagrange Inversion). *For any function of the form $y = x\varphi(y)$, where φ is a power series such that $\varphi(0) = 1$,*

$$[x^n]\psi(y) = \frac{1}{n}[u^{n-1}]\varphi(u)^n\psi'(u).$$

In particular, for $\psi(y) = y^k$, one has

$$[x^n]y^k = \frac{k}{n}[u^{n-k}](1-u)^{-n}.$$

Proof of Theorem 3.2. The first parts are easy to see. Then, one first shows that $|Q_r|$ is given by the coefficient of x^r in the expression $c(x)^{n-r-1}$, where $c(x)$ is the generating function for the Catalan numbers, $c(x) = 1 + C_1x + C_2x^2 + \dots = 1 + x + 2x^2 + 5x^3 + 14x^4 + \dots$.

Fix a tree $T \in Q_r$. Note that from Corollary 2.11(v), the length of the right arm of T is $n - r - 1$, and so there are precisely $n - r - 1$ vertices on the right arm of T which have (possibly single-leaf) left subtrees. The total number of leaves on these left subtrees is $n - 1$. So, we are looking for the number of ways of distributing $n - 1$ leaves among $n - r - 1$ “blank” subtrees, such that the subtrees are also binary trees in their own right. We can simplify the problem by putting a leaf in each subtree to begin with. Thus, we are looking for the number of ways of putting $n - 1 - (n - r - 1) = r$ indistinguishable objects in $n - r - 1$ numbered boxes, with each box containing i objects weighted C_i .

The number of ways of doing this is given by the coefficient of x^r in $(1 + C_1x + C_2x^2 + \dots)^{n-r-1}$, i.e. $c(x)^{n-r-1}$, where $c(x)$ is as above.

To complete the proof, use Lagrange Inversion as follows: if $c(x)$ is defined as above, then set $y(x) = xc(x)$. Then, y satisfies $y = x(1 - y)^{-1}$. Now, using Lagrange Inversion from Lemma 3.3 above with $k = n - r - 1$ and $m = n - 1$, one has, after a few simplifications,

$$|Q_r| = \frac{n - r - 1}{n - 1} \binom{n + r - 2}{r} = \binom{n + r - 2}{r} - \binom{n + r - 2}{r - 1}. \quad \square$$

Corollary 3.4. *The sizes of the ranks in $\mathcal{C}_n$ weakly increase. In fact, they strictly increase until the final rank Q_{n-2} , which has the same size, C_{n-2} , as the penultimate rank Q_{n-3} .*

Proof. From Theorem 3.2, it can be seen that $|Q_i| = \frac{(n+i-2)!}{i!(n-1)!} \cdot (n - i - 1)$, and so, for consecutive ranks r and $r + 1$, one has

$$\frac{|Q_{r+1}|}{|Q_r|} = \frac{(n + r - 1)(n - 2 - r)}{(r + 1)(n - 1 - r)}.$$

The rank size increases weakly whenever the numerator is at least as large as the denominator, and hence the condition for weakly increasing rank size is $(n + r - 1)(n - r - 2) \geq (r + 1)(n - r - 1)$. But this condition reduces after a few simple manipulations to the condition $n^2 - 4n + 3 - r(n - 1) \geq 0$. The result can be verified easily. $\square$

4. DISTANCES IN $\mathcal{C}_n$ AND $\mathcal{R}_n$

We now prove some properties of the comb poset relating to the distance between pairs of trees in the rotation graph $\mathcal{R}_n$.

Proposition 4.1. *Any ascending chain in the right comb poset $\mathcal{C}_n$ is an ascending chain in $\mathcal{T}_n$.*

Proof. From Corollary 2.11(ii), one has that T_2 is a cover of T_1 in $\mathcal{C}_n$ iff RP_{T_2} can be obtained from RP_{T_1} by adding precisely one more parenthesis pair. Adding any parenthesis pair to RP_T is the same as shifting a pair of parentheses to the *left* in the corresponding full parenthesization of the leaves of T . $\square$

Proposition 4.2. *Suppose T_1 and T_2 are two trees have a common upper bound in $\mathcal{C}_n$. Furthermore, suppose there are pairs of parentheses J_1 in RP_{T_1} and J_2 in RP_{T_2} such that J_1 and J_2 enclose a common factor (which may be a left factor for one and a right one for the other). Then, $J_1 = J_2$.*

Proof. Suppose first that the common factor E is a left factor in J_1 and a right factor in J_2 . Then, because $T_1 \vee T_2$ exists, from Corollary 2.11(i), one has that J_1 and J_2 must be in $RP_{T_1 \vee T_2}$ as well, which then implies that E is enclosed as a single factor by a pair of parentheses in $RP_{T_1 \vee T_2}$, contradicting Proposition 2.2. So, assume without loss of generality that E is the *left* factor for both J_1 and J_2 . If we denote the right factors of J_1 and J_2 by E_1 and E_2 , respectively, then these must both be the right factor of the corresponding parenthesis pair in $RP_{T_1 \vee T_2}$, forcing $E_1 = E_2$, and hence $J_1 = J_2$. $\square$

Theorem 4.3. *If T_1 and T_2 are two trees in some interval in $\mathcal{C}_n$, then the shortest distance between them along the edges of the rotation graph $\mathcal{R}_n$ is given by*

$$d_{\mathcal{R}_n}(T_1, T_2) = \text{rank}(T_1) + \text{rank}(T_2) - 2 \cdot \text{rank}(T_1 \wedge T_2).$$

Equivalently, from Corollary 2.11(iv), this shortest distance is also given by

$$d_{\mathcal{R}_n}(T_1, T_2) = 2 \cdot \text{rank}(T_1 \vee T_2) - \text{rank}(T_1) - \text{rank}(T_2).$$

Proof. From Corollary 2.11(i) one has that $RP_{T_1 \wedge T_2}$ contains all the common parenthesis pairs of RP_{T_1} and RP_{T_2} . Hence, RP_{T_1} and RP_{T_2} are formed by adding respectively some r and s *extra* pairs of parentheses to $RP_{T_1 \wedge T_2}$, from Corollary 2.11(ii), where r and s are nonnegative integers.

Suppose $(\lambda_1, \lambda_2, \dots, \lambda_\ell)$ is a path from T_1 to T_2 in $\mathcal{R}_n$, with λ_i being the rotation between trees, say, S_i and S_{i+1} , with λ_ℓ the rotation from S_ℓ to T_2 . Define a map f from $RP_{T_1} \triangle RP_{T_2}$, the symmetric difference of the sets of parentheses in RP_{T_1} and RP_{T_2} , to $\{\lambda_1, \lambda_2, \dots, \lambda_\ell\}$ by $f(J) = \lambda_j$

where j is the minimum index such that λ_j impacts J , so $S_1 = T_1$. This map will be shown to be injective.

Suppose $f(J) = f(K) = \lambda_j$. If $J \neq K$, one may assume without loss of generality that λ_j is the rotation sending J to K . Then J and K together enclose factors A, B, C arranged as $(A(B)C)$. While λ_j is not necessarily a left rotation centered on the right arm (since λ_j represents an arbitrary edge in $\mathcal{R}_n$), it is centered at the root of some subtree of S_j , call it S . Then λ_j is a rotation sending S to S' , which is itself a subtree of S_{j+1} having K as a parenthesis pair. In S , λ_j is a right arm rotation, so S' covers S in $\mathcal{C}_m$ for some $m \leq n$. Then Proposition 4.2 forces $J = K$, a contradiction since λ_j was a rotation moving J to K .

Thus the map f is injective and the minimum length of a path is $|RP_{T_1} \triangle RP_{T_2}| = \text{rank}(T_1) + \text{rank}(T_2) - 2 \cdot \text{rank}(T_1 \wedge T_2)$. Furthermore, we know a path of this length exists – the path in $\mathcal{C}_n$ from T_1 to $T_1 \wedge T_2$ by deleting the r extra pairs of parentheses in RP_{T_1} , and then from $T_1 \wedge T_2$ to T_2 by adding in the s extra pairs in RP_{T_2} , proving the result. $\square$

Corollary 4.4. *For trees T_1 and T_2 with an upper bound in $\mathcal{C}_n$, if RP_{T_1} and RP_{T_2} are obtained by adding some r extra pairs $\{J_i\}$ and s extra pairs $\{J'_i\}$ of parentheses to $RP_{T_1 \wedge T_2}$, then*

- (i) J_i and J'_j are disjoint for all $1 \leq i \leq r$ and $1 \leq j \leq s$.
- (ii) Any shortest path between T_1 and T_2 in $\mathcal{R}_n$ has length $r + s$.
- (iii) In any such shortest path, precisely r moves correspond to removing or moving the $\{J_i\}$, and precisely s moves correspond to adding the $\{J'_i\}$, and no move can achieve both simultaneously.

Conjecture 4.5. *For T_1, T_2 with an upper bound in $\mathcal{C}_n$, a shortest path in $\mathcal{R}_n$ from T_1 to T_2 also lies in $\mathcal{C}_n$.*

This amounts to proving that for any pair of trees T_1 and T_2 with an upper bound in $\mathcal{C}_n$, any shortest path between them in $\mathcal{R}_n$ consists of simply adding and deleting parenthesis pairs from their RP_T 's, and not of any “shifting” move. The conjecture appears to be true, but the authors have not been able to find a satisfactory argument.

Corollary 4.6. *The rank of any tree $T \in \mathbb{T}_n$ in $\mathcal{C}_n$ is its distance from the right comb tree along the edges of the rotation graph $\mathcal{R}_n$. Furthermore, from Corollary 2.11(iii), the distance of T from the right comb tree in $\mathcal{R}_n$ is given by the number of parenthesis pairs in RP_T .*

Remark. It can be easily shown from the result above that the *diameter* of the rotation graph $\mathcal{R}_n$, given by the maximum distance between any pair of trees in $\mathcal{R}_n$, is at most $2n - 4$ for any $n \in \mathbb{N}$. In [8], Sleator, Tarjan and Thurston established tighter bounds on the diameter of the rotation graph.

5. TAMARI MEETS AND JOINS FOR TWO TREES IN SOME INTERVAL

From Corollary 2.11(i), we know the meaning of the meet and join of a pair of trees having a common upper bound in our poset. It is natural to ask how these meets and joins relate to meets and joins in the Tamari lattice. As before, we will refer to meets and joins in the Tamari lattice $\mathcal{T}_n$ as the “Tamari meet” and “Tamari join”.

The first observation is that, while two arbitrary trees in $\mathcal{C}_n$ do have a well-defined meet in $\mathcal{C}_n$, this meet does not necessarily correspond to the Tamari meet. For example, consider the pair of trees represented by $T_1 = (((a_1 a_2) a_3) a_4) a_5$ and $T_2 = ((a_1 (a_2 a_3)) a_4) a_5$. This pair has Tamari meet T_2 , while their meet in $\mathcal{C}_n$ is just the right comb tree. Further, recall that $\mathcal{C}_n$ is a meet-semilattice rather than a lattice, so not all pairs of trees have a join.

However, something much stronger can be said if both the trees under consideration are in some interval in the comb poset; it turns out that their meet and join in $\mathcal{C}_n$ correspond to their Tamari meet and join.

Recall the concept of pruned trees from Definition 2.7. Furthermore, there is a well-known natural numbering of the vertices of the pruned tree using $1, 2, \dots, n - 1$, in which a vertex receives a higher number than any vertex in its left subtree, but a lower one than any vertex in its right subtree. This labeling is unique, and well known. It is called the *in-order* labeling of a pruned tree on $n - 1$ vertices.

Example. Figure 4 shows a pruned tree on 8 vertices, corresponding to the 9-leaf binary tree whose reduced parenthesization is $((a_1a_2)(a_3a_4))a_5(a_6(a_7a_8))a_9$, is labeled in the in-order labeling.




FIGURE 4.

The subsequent discussion uses the notion of *bracketing vectors*. A more formal treatment than we will give here can be found in Huang and Tamari's original paper, [3].

Definition 5.1. Consider the “pruned” binary tree representation of some tree $T \in \mathbb{T}_n$, and number the $n - 1$ vertices by the in-order labeling. Then, the *bracketing vector* for T , $\langle T \rangle = \langle b_1(T), \dots, b_{n-1}(T) \rangle$, has $b_j(T)$ equal to the number of vertices in the left subtree of the vertex labeled j in the pruned tree. In particular, the first coordinate of a bracketing vector is always 0.

Example. The bracketing vector for the tree in Figure 4 is the 8-tuple $(0, 1, 0, 3, 0, 0, 0, 2)$.

For any $T \in \mathbb{T}_n$, take its corresponding RP_T , and then enclose every leaf by an additional parenthesis pair (which therefore encloses just one factor, the leaf itself). Then, for $i \in \{1, \dots, n-1\}$, take the largest parenthesis pair such that the leaf a_i is the largest leaf enclosed by that parenthesis pair in the representation above. Denote the number of leaves enclosed by this largest parenthesis pair by k_i and let $k = (k_1, \dots, k_{n-1})$.

Proposition 5.2. For $T \in \mathbb{T}_n$, the i^{th} component of the bracketing vector for T is precisely $k_i - 1$.

Proof. From Proposition 2.9, one has that the unbracketed elements in RP_T correspond to vertices on the right arm that have a single leaf rather than a left subtree (in other words, their coordinate in the bracketing vector is 0). This gives the desired result for $k_i = 1$. For $k_i > 1$, we can follow a recursive argument by considering the bracketed elements as trees in their own right. $\square$

Example. Consider the tree T having reduced parenthesization $((a_1a_2)(a_3a_4))a_5(a_6(a_7a_8))a_9$. Adding the additional parentheses gives

$$(((a_1)(a_2))((a_3)(a_4)))(a_5)((a_6)((a_7)(a_8)))(a_9).$$

Then $k = (1, 2, 1, 4, 1, 1, 3)$ and $\langle T \rangle = (0, 1, 0, 3, 0, 0, 0, 2)$.

The connection between bracketing vectors and meets and joins in the Tamari lattice rests on a result of Pallo. The proof may be found in [6, Theorem 2]. The k -vector defined above is Pallo's weight vector w_T .

Theorem 5.3 (Pallo). *For two n -leaf binary trees T and T' , one has $T \leq T'$ if and only if the bracketing vector of T is component-wise less than or equal to the bracketing vector of T' . Furthermore, the bracketing vector for the meet of two trees in the Tamari lattice corresponds to the componentwise minimum of the bracketing vectors of the two trees.*

Theorem 5.4. *Let $\langle T \rangle$ denote the bracketing vector for T . Let T_1 and T_2 be arbitrary trees in the same interval of $\mathcal{C}_n$. Then, their meet and join in $\mathcal{C}_n$ are given by the trees corresponding respectively to the componentwise minimum and the componentwise maximum of $\langle T_1 \rangle$ and $\langle T_2 \rangle$.*

Proof. For any coordinate i of $\langle T_1 \vee T_2 \rangle$, the corresponding leaf a_i is the largest leaf in one or more parenthesis pairs in the representation described above, before Proposition 5.2. From Corollary 2.11(i), one has that because these parenthesis pairs represent the union of pairs from RP_{T_1} and RP_{T_2} , the same parenthesis pairs must be present in at least one of RP_{T_1} and RP_{T_2} . Furthermore,

any *bigger* pair in RP_{T_1} or RP_{T_2} that has a_i as the largest enclosed leaf would also have to appear in $RP_{T_1 \vee T_2}$, a contradiction. From Proposition 5.2 above, this proves the theorem for joins, and the proof for meets is analogous. $\square$

Corollary 5.5. *For T_1 and T_2 in some interval in $\mathcal{C}_n$, their meet and join in $\mathcal{C}_n$ correspond respectively to their meet and join in the Tamari lattice $\mathcal{T}_n$.*

Proof. The proof for the meet follows directly from Theorems 5.3 and 5.4. The proof for the join is obtained by the same argument that Pallo uses in Theorem 2 of [6]. $\square$

6. RELATION WITH A POSET OF EDELMAN

In [2], Edelman introduced a subposet of the right weak order on the symmetric group $\mathfrak{S}_n$. Although this poset is not a lattice, the intervals are known to each be distributive lattices, as is the case for the comb poset $\mathcal{C}_n$.

Definition 6.1. The *right weak order* on $\mathfrak{S}_n$ is a partial ordering of the elements of $\mathfrak{S}_n$ defined as the transitive closure of the following covering relation: a permutation σ covers a permutation τ if σ is obtained from τ by a transposition of two adjacent elements of the one line notation of τ introducing an inversion.

Edelman imposed an additional constraint on this ordering, under which σ covers τ , if, after the transposition of x_j and x_{j+1} as above, nothing to the left of x_{j+1} in σ is greater than x_{j+1} . This restriction results in a subposet of the right weak ordering on $\mathfrak{S}_n$. Denote this poset by $\mathcal{E}_n$.

Example. Figure 5 depicts the Hasse diagram of $\mathcal{E}_3$, with an additional dashed edge indicating the extra order relation in the right weak order on $\mathfrak{S}_3$.




FIGURE 5. Edelman's Poset $\mathcal{E}_3$.

Definition 6.2. The *pruned tree map*, $p: \mathfrak{S}_n \rightarrow \{\text{pruned trees on } n \text{ vertices}\}$, is defined recursively as follows. For $x \in \mathfrak{S}_1$, $p(x)$ is the tree with a single vertex. Then, for $n > 1$ and $x \in \mathfrak{S}_n$, define

$$p(x) = \begin{array}{c} \bullet \\ \swarrow \quad \searrow \\ \bullet \qquad \bullet \\ p(x_{<}) \quad p(x_{>}) \end{array}$$

where $x_{<} = (x_{i_1}, \dots, x_{i_k})$ where $i_1 < \dots < i_k$ are the indices of all elements of x less than x_1 and $x_{>}$ is defined similarly for elements of x greater than x_1 . Extend p to a map $\beta: \mathfrak{S}_n \rightarrow \mathbb{T}_{n+1}$ by attaching leaves to $p(x)$ to give a binary tree (in other words, “unpruning” $p(x)$).

Remark. Amending the definition of p slightly so that the root of $p(x)$ is labeled by x_1 results in the pruned tree having the in-order labeling (see Section 5). This labeled tree is, in fact, the unbalanced binary search tree for the permutation. (See [5].) The pruned tree map is also related to the bijection between permutations and increasing binary trees on n vertices (see [9, p. 24]): the pruned tree associated to w is the increasing binary tree associated to w^{-1} with the labels removed. Consequently, the pruned tree map is a surjection.

Example. Figure 6 shows $p: \mathfrak{S}_4 \rightarrow \{\text{pruned trees with 4 vertices}\}$. Permutations having the same image are circled.




FIGURE 6.

Theorem 6.3. *The map $p: \mathfrak{S}_n \rightarrow \mathbb{T}_{n+1}$ gives an order-preserving surjection from $\mathcal{E}_n$ to $\mathcal{C}_{n+1}$.*

Proof. As noted above, p is a surjection. It suffices to show that if τ covers σ in $\mathcal{E}_n$ and T_1 and T_2 are the images of σ and τ , respectively, under p , then either $T_1 = T_2$, or T_2 covers T_1 in $\mathcal{C}_{n+1}$.


Suppose

$$\sigma = (x_1, x_2, \dots, x_j, x_{j+1}, \dots, x_n) \in \mathfrak{S}_n \quad \text{and} \quad \tau = (x_1, x_2, \dots, x_{j+1}, x_j, x_{j+2}, \dots, x_n) \in \mathfrak{S}_n,$$

with τ covering σ in $\mathcal{E}_n$. From the definition of a cover in $\mathcal{E}_n$ one has $x_j < x_{j+1}$. Now, if $j = 1$, then this transposition corresponds to a left rotation centered on the root, and therefore T_2 covers T_1 in $\mathcal{C}_{n+1}$. So assume $j \neq 1$; in other words, x_j is not the root x_1 of the tree.


Since τ covers σ in $\mathcal{E}_n$, one knows that $x_j > x_s$ for all $s < j$. Suppose there is $s < j$ such that $x_{j+1} < x_s < x_j$. Then, by the definition of the pruned tree map, the vertex labeled by x_{j+1} lies in the left subtree of the vertex labeled by x_s and x_j lies in the right subtree of the image of σ . However, this is also true of the image of τ . Since the transposition of x_j and x_{j+1} is the only difference between σ and τ , we must have $T_1 = T_2$.

Consequently, if $T_1 \neq T_2$, one may assume there is no $s < j$ with $x_{j+1} < x_s < x_j$. In such a case, T_1 must be of the form



Here the white circle S denotes the parent tree of the entire subtree shown, with the condition that x_j and x_{j+1} lie on the right arm. The white circles X , Y and Z denote arbitrary subtrees, whose interpretations in terms of the elements in σ are as follows: X is the image under P of the ordered sequence of elements appearing *after* x_j which are less than x_j , while Z is the ordered sequence of elements appearing *after* x_{j+1} which are greater than x_{j+1} , and Y is the ordered sequence of elements appearing after x_j that lie between x_j and x_{j+1} .

Now, consider what happens to T_2 , when x_j and x_{j+1} are switched. The tree T_2 is depicted below.




Here, S is going to be unchanged, and x_j and x_{j+1} must move as shown. In addition, there will be subtrees X' , Y' , and Z' as drawn above. However, notice that, if one considers what these subtrees must be with respect to the permutation τ , the fact that x_j and x_{j+1} are adjacent forces the conclusion that the subtrees are unchanged from σ , or in other words that $X = X'$, $Y = Y'$ and $Z = Z'$. So then, T_2 is obtained by a left rotation centered on a vertex on the right arm of T_1 . Therefore, T_2 covers T_1 in $\mathcal{E}_{n+1}$, and hence we are done. $\square$

To relate the intervals of $\mathcal{E}_{n+1}$ to those of $\mathcal{E}_n$ more deeply, a formal discussion of $\mathcal{E}_n$ is needed. In [2], Edelman defined the following order on the inversion set of a permutation σ .

Definition 6.4. Define $I(\sigma) := \{(j, i) : j > i \text{ and } \sigma^{-1}(j) < \sigma^{-1}(i)\}$. Order $I(\sigma)$, with $(k, \ell) \geq (j, i)$ if and only if $k \geq j$ and $\sigma^{-1}(\ell) \leq \sigma^{-1}(i)$. In a slight abuse of notation, the poset $(I(\sigma), <)$ shall be referred to as $I(\sigma)$ as well.

Theorem 6.5 (Edelman, [2, Theorem 2.13]). $[e, w]_{\mathcal{E}_n} \simeq J(I(w))$, where $[e, w]_{\mathcal{E}_n} = \{v \in \mathfrak{S}_n : v \leq_{\mathcal{E}_n} w\}$, via $v \mapsto I(v)$.

Definition 6.6. Fix a permutation $w \in \mathfrak{S}_n$. Let T_w be the image of w under the pruned tree map, p . Recall the reduced pruned poset from Definition 2.8. Here it will be useful to label its vertices by the labels they have in T_w , rather than by parentheses. Define a map $f : P_{T_w} \rightarrow I(w)$ as follows: $f(j) = (i, j)$, where i is the smallest label of a vertex of T_w such that j lies in the left subtree of i .

FIGURE 7. T_w , P_{T_w} and $I(w)$ with the image of f indicated.

Example. Suppose $w = (4, 9, 2, 1, 8, 3, 6, 7, 5) \in \mathfrak{S}_9$. Figure 7 depicts T_w , P_{T_w} and $I(w)$, with the image of f indicated in $I(w)$.


Proposition 6.7. *The map f is order-preserving.*

Proof. It suffices to show that if $j > k$, and j covers k in P_{T_w} , then $f(j) > f(k)$. Since j covers k , one has that k is a child of j , and there are two cases.

- (1) If k is a left child of j , then $f(k) = (j, k)$. By the definition of the pruned tree map (Definition 6.2), one knows $w^{-1}(j) < w^{-1}(k)$. Suppose $f(j) = (i, j)$. Then, by definition, $j < i$, which means that $(j, k) < (i, j)$ in $I(w)$, as desired.
- (2) If k is a right child of j , then $f(j) = (i, j)$ means that $f(k) = (i, k)$. Now, $w^{-1}(j) < w^{-1}(k)$, and so $(i, j) > (i, k)$, as desired.

These cover all the cases, proving the result. $\square$

Definition 6.8. Let P_1, P_2 be two posets and suppose $\phi: P_1 \rightarrow P_2$ is order-preserving. Then ϕ induces a map $J(\phi): J(P_2) \rightarrow J(P_1)$ defined by $J(\phi)(I) = \phi^{-1}(I)$. One calls $J(\phi)$ the *Birkhoff-Priestley dual* to f . In fact, $J(\phi)$ is a lattice morphism.

FIGURE 8. $p|_{[e, 4213]}$ and $J(f)$ for the interval $[e, 4213]_{\mathcal{E}_n}$.

Remark. The ensuing discussion will make use of the following observation. Given an order ideal in P_{T_w} , associated to it is an in-order labeled pruned tree.

- Each connected component of the Hasse diagram will be the left subtree of some element on the right arm of the pruned tree.
- Any elements of $[n]$ not in L_T are arranged as the right arm of the tree, in descending order.
- Given a connected component containing $\{i_1, \dots, i_k\}$, it is the left subtree of $\max\{i_j\} + 1$, where we take the maximum in numerical order, and not the P_{T_w} order.

Theorem 6.9. *For each $w \in \mathfrak{S}_n$, the map $f: P_{T_w} \rightarrow I(w)$ defined in Definition 6.6 is Birkhoff-Priestley dual to the pruned tree map $p: [e, w]_{\mathcal{E}_n} \rightarrow J(P_{T_w})$. In particular, $p: \mathcal{E}_n \rightarrow \mathcal{C}_n$ becomes a lattice morphism when restricted to any interval in $\mathcal{E}_n$. As a commutative diagram, one has*

$$\begin{array}{ccc}
 \mathcal{E}_n & \xrightarrow{p} & \mathcal{C}_{n+1} \\
 \uparrow & & \uparrow \\
 [e, w]_{\mathcal{E}_n} & \xrightarrow{p|_{[e, w]_{\mathcal{E}_n}}} & [\text{RCT}(n+1), T_w]_{\mathcal{C}_{n+1}} \\
 \uparrow \wr & & \uparrow \wr \\
 J(I(w)) & \xrightarrow{J(f)} & J(P_{T_w})
 \end{array}$$

Example. Figure 8 depicts Theorem 6.9 on the interval $[e, 4213]_{\mathcal{E}_n}$.

Proof. Begin by noting that, strictly speaking, the Birkhoff-Priestley dual to f , $J(f)$, is not a map from $[e, w]_{\mathcal{E}_n} \rightarrow J(P_{T_w})$ as p is, but $J(f): J(I(w)) \rightarrow J(P_{T_w})$. However, from Theorem 6.5, $J(I(w)) \simeq [e, w]_{\mathcal{E}_n}$, so one can use p in place of such a $J(f)$.


Fix $w \in \mathfrak{S}_n$. From Theorem 6.3 one has that $p: [e, w]_{\mathcal{E}_n} \rightarrow J(P_{T_w})$ is order-preserving. Then one must show that $p(I)$ is, in fact, $f^{-1}(I)$.

Induct on the number of inversions in a permutation in $[e, w]_{\mathcal{E}_n} \simeq J(I(w))$. Note that the claim is trivially true for $(1, 2, \dots, n)$, the identity permutation, which corresponds to $\emptyset \in J(I(w))$.


Now suppose the permutation $\tau \in [e, w]_{\mathcal{E}_n}$, and $f^{-1}(I(\tau)) = T_\tau$. Suppose σ covers τ . Then, σ has precisely one more inversion than τ ; call this pair (i, j) . There are, again, two cases:

- (1) If (i, j) is not in the image of f , then $f^{-1}(I(\sigma)) = f^{-1}(I(\tau))$, and so one must show that $T_\sigma = T_\tau$. We know i and j are adjacent in τ , and i is a left-right maximum. In particular, this means that neither τ nor σ has an inversion (k, i) , meaning i lies in the right arm of both T_τ and T_σ . Again, there are two subcases:

- Suppose there is an inversion (ℓ, j) with $\tau^{-1}(\ell) = \sigma^{-1}(\ell) < \sigma^{-1}(i)$. Then, j is in the left subtree of ℓ in T_τ and T_σ , meaning that it is not the parents of i in T_τ , and so adding the inversion (i, j) does not change T_τ , forcing $T_\tau = T_\sigma$, as desired.
- Now suppose there is no “earlier” inversion, and j is a left-right maximum. Then one has




Since (i, j) is not in the image of f , there must be some “later” inversion moving k past j , changing a permutation α to a permutation β on the way to w , such that the addition of this inversion changes the image under the pruned tree map. In other words, k must be the right child of j in T_α , giving




This means that k was in C , as i 's left subtree in T_τ is labeled, and so $k < i$. This means that in σ one has $(\dots, i, j, \dots, k)$, and so k cannot possibly be a left-right maximum in α , to move past j to obtain β , a contradiction. Hence, one must have $f(j) = (i, j)$, contradicting the assumption that (i, j) was not in the image of f .

- (2) In the second case, suppose (i, j) is in the image of f . Then, the addition of the inversion of (i, j) to τ results in σ , and in the following rotation from T_τ to T_σ .



One needs to show that $f^{-1}(I(\sigma))$ is the order ideal in P_{T_w} that “attaches” to form T_σ . Now, since T_τ and T_σ differ only in this rotation, one need only show that



appears in P_{T_w} . Left-right maxima occur only on the right arm of the image of a permutation under the pruned tree map, and so subsequent inversions on the way from σ to w result in rotations in the pruned tree that cannot affect the children of j . Hence, the above subtree appears in P_{T_w} , and so, T_σ is the pruned tree associated to $I(\sigma)$.

These two cases cover all possibilities, concluding the proof. $\square$

7. THE PARSEWORDS FUNCTION FOR THE COMB POSET

The number of common parsewords for any two trees having a common upper bound in $\mathcal{C}_n$ can be computed precisely. Recall that $w \in \text{ParseWords}(T)$ means that T admits a labeling of its vertices by $0, 1, 2$ such that the leaves are labeled by the word w , the children of each vertex have distinct labels and no vertex has the same label as either of its children. Kauffman showed in [4] that the Four Color Theorem is equivalent to $\text{ParseWords}(T_1, T_2) \neq \emptyset$ for all $T_1, T_2 \in \mathbb{T}_n$ for any $n \in \mathbb{N}$. It was further recent work by Cooper, Rowland and Zeilberger in [1] that led us to first consider the comb poset.

Example. An example of a tree parsing the word 2202 is shown in Figure 9.




FIGURE 9.

Example. An example of two trees parsing the same word 010 is shown in Figure 10.




FIGURE 10.

Example. The common parsewords for the trees in Figure 10 are 101, 202, 010, 212, 020, 121.


To simplify notation, let $T_{\leq b}$ be the subtree of a tree T having the vertex b as its root.

Proposition 7.1 (Common root property, [1, Proposition 2]). *If two trees $T_1, T_2 \in \mathbb{T}_n$ parse the same word, then their roots receive the same label when the trees are labeled with a common parse-word. Hence, if for $T_1, T_2 \in \mathbb{T}_n$, there are vertices b_i in T_1 and b_j in T_2 such that $T_{1 \leq b_i}$ and $T_{2 \leq b_j}$ have precisely the same leaves (i.e. both the dangling subtrees contain precisely the leaves m_1 through m_2 , for some natural numbers $m_1 < m_2 \leq n$), then b_i and b_j receive the same label if we label the trees with a common parse word.*


If a tree T parses a word w and X is a subtree of T , let $w(X)$ be the label received by the root of X parsing w and let w_X be the segment of w parsed by the subtree X .

Definition 7.2. Say $T \in \mathbb{T}_n$ has a *leaf reduction* at $(i, i+1)$ for $i \in \{0, 1, \dots, n\}$ if


- (i) the leaves $i, i+1$ have a common parent:



- (ii) the leaf ℓ_1 is attached to the root r if $i = 0$:



- (iii) the leaf ℓ_n is attached to the root r if $i = n$:



Define $\hat{T}$ as in the above diagrams, i.e. remove either ℓ_i, ℓ_{i+1} , ℓ_1 or ℓ_n from T , depending on the case, and define two maps $f_i^<, f_i^>: \text{ParseWords}(\hat{T}) \rightarrow \text{ParseWords}(T)$ sending $\hat{w}$ to the w in $\text{ParseWords}(T)$ that uniquely extends $\hat{w}$ in such a way that $w_i < w_{i+1}$ or $w_i > w_{i+1}$, respectively. (If $i = 0$ or $i = n$, treat w_{i+1} and w_i , respectively, as $w(\hat{T})$.)

Proposition 7.3. *If $\{T_1, \dots, T_m\} \subset \mathbb{T}_n$ share a leaf reduction at $(i, i+1)$, then*

$$\text{ParseWords}(T_1, \dots, T_m) = f_i^<(\text{ParseWords}(\hat{T}_1, \dots, \hat{T}_m)) \sqcup f_i^>(\text{ParseWords}(\hat{T}_1, \dots, \hat{T}_m)).$$

Proposition 7.3 is most frequently used several times in succession, to “collapse” a subtree common to two or more trees. In particular, it often allows a reduction to the case $T_1 \wedge T_2 \wedge \dots \wedge T_m = \text{RCT}(n)$.

Corollary 7.4. *For $T \in \mathbb{T}_n$, one has $|\text{ParseWords}(T)| = 3 \cdot 2^{n-1}$.*


Proof. The case $n = 1$ is trivial. One can then induct on n , applying Proposition 7.3 to T alone. $\square$

Proposition 7.5. *For any $T_1, T_2 \in \mathbb{T}_n$ differing by a single rotation (not necessarily a right arm rotation),*

$$\begin{aligned} \text{ParseWords}(T_1, T_2) &= \{w \in \text{ParseWords}(T_1) : w(X) \neq w(Y)\} \\ &= \{w \in \text{ParseWords}(T_1) : w(Y) \neq w(Z)\} \\ &= \{w \in \text{ParseWords}(T_1) : w(X) = w(Z)\}, \end{aligned}$$

where X, Y, Z are subtrees as indicated below. Furthermore, $|\text{ParseWords}(T_1, T_2)| = 3 \cdot 2^{n-2}$.

Proof. The conditions on w in the first part of the claim can be checked by inspection. For the second part, the case $n = 3$ can be checked directly. Taking this as a base case, one can induct on n . A rotation looks like




Applying Proposition 7.3 to any of the subtrees X, Y, Z not consisting of a single leaf, or to the subtrees taken together as a single subtree if all three are leaves, allows one to invoke the result for a smaller n and obtain the desired result. $\square$

Theorem 7.6. *Suppose $T_1 < T < T_2$ in $\mathcal{C}_n$. Then $\text{ParseWords}(T_1, T_2) = \text{ParseWords}(T_1, T_2, T)$.*

Proof. It suffices to prove the statement for $T_1 < T < T_2$ in $\mathcal{C}_n$. Assume the theorem holds in this case and obtain the general case by induction on the length of a chain between T_1 and T .

Suppose we have $T_1 < T'_1 < T'_2 < \dots < T'_k < T < T_2$. Then, by induction, $\text{ParseWords}(T_1, T'_k, T_2) = \text{ParseWords}(T_1, T_2)$. Suppose $w \in \text{ParseWords}(T_1, T_2) = \text{ParseWords}(T_1, T'_k, T_2)$. Furthermore, $w \in \text{ParseWords}(T'_k, T_2) = \text{ParseWords}(T'_k, T, T_2)$, so $w \in \text{ParseWords}(T_1, T, T_2)$ as desired. By definition, $\text{ParseWords}(T_1, T, T_2) \subset \text{ParseWords}(T_1, T_2)$, so $\text{ParseWords}(T_1, T, T_2) = \text{ParseWords}(T_1, T_2)$, as desired.

To prove the initial case, now suppose $T_1 < T < T_2$. One has a sequence of right-arm rotations



Since the rotation between T_1 and T moves the subtrees labeled by X and Y off the right arm, they must remain in the same position relative to one another in T_2 .


Suppose $w \in \text{ParseWords}(T_1, T_2)$. Since w parses T_2 , one must have $w(X) \neq w(Y)$ and, hence, by Proposition 7.5, w parses T . Thus $\text{ParseWords}(T_1, T_2) \subset \text{ParseWords}(T_1, T, T_2)$. By definition, $\text{ParseWords}(T_1, T, T_2) \subset \text{ParseWords}(T_1, T_2)$, so $\text{ParseWords}(T_1, T_2) = \text{ParseWords}(T_1, T, T_2)$, as desired. $\square$

Theorem 7.7. *Suppose $T < T'$ in $\mathcal{C}_n$ and $\text{rank}(T') - \text{rank}(T) = k$. Then $|\text{ParseWords}(T, T')| = 3 \cdot 2^{n-1-k}$.*

Proof. One proceeds by induction. Proposition 7.5 addresses the case $k = 1$. Via repeated leaf reductions, one may assume $T = \text{RCT}(n)$. Now suppose the statement holds for $k - 1$, that $T < T'$ and $\text{rank}(T') - \text{rank}(T) = k$. One has a chain in $\mathcal{C}_n$, $T < T_1 < T_2 < \dots < T_{k-1} < T'$. By induction $|\text{ParseWords}(T, T_{k-1})| = 3 \cdot 2^{n-k}$. We will construct a bijection

$$\text{ParseWords}(T, T_{k-1}, T') \rightarrow \text{ParseWords}(T, T_{k-1}, T') \setminus \text{ParseWords}(T, T_{k-1}, T').$$

First, one characterizes those parsewords in $\text{ParseWords}(T, T_{k-1}, T')$. Since T' covers T_{k-1} , one has that they differ by a right arm rotation:




From Proposition 7.5, one has that $w \in \text{ParseWords}(T, T_{k-1})$ also parses T' (i.e. $w \in \text{ParseWords}(T, T_{k-1}, T')$) if and only if $w(X) \neq w(Y)$.

Now define the map


$$\phi: \text{ParseWords}(T, T_{k-1}) \rightarrow \text{ParseWords}(T, T_{k-1})$$

as follows. Suppose $w \in \text{ParseWords}(T, T_{k-1})$. Then $w = w_S w_X w_Y w_Z$, where w_J is the word parsing the leaves of the subtree J . Define a transposition in $\mathfrak{S}_{\{0,1,2\}}$ by $\sigma = (w(Y), w(Z))$. Then define $\phi(w) = w_S w_X \sigma(w_Y) \sigma(w_Z)$.

On the one hand, ϕ permutes the alphabet with the subtree of T_{k-1} consisting of Y and Z , while leaving the label of its root the same, so $\phi(w)$ certainly parses T_{k-1} . Recall that T was assumed to be $\text{RCT}(n)$. Labeling T with w gives



Fully labeling the subtree of T containing the leaves of T_{k-1} 's Y and Z subtrees entails giving the root the label $w(Y)$ and another right arm vertex the label $w(Z)$:



□

Then applying σ to the vertices of T in the Y and Z subtrees of T_{k-1} separately is the same as applying σ to both the Y and Z parts of T simultaneously. Thus $\phi(w)$ parses T , so the map is well-defined.

Then ϕ is transparently a bijection

$$\text{ParseWords}(T, T_{k-1}) \rightarrow \text{ParseWords}(T, T_{k-1})$$

and, moreover, exchanges $\text{ParseWords}(T, T_{k-1}, T')$ and $\text{ParseWords}(T, T_{k-1}) \setminus \text{ParseWords}(T, T_{k-1}, T')$: if $w \in \text{ParseWords}(T, T_{k-1}, T')$, then $\phi(w)$ has $\phi(w)(X) = \phi(W)(Y)$, meaning it cannot parse T' . Thus, ϕ is a bijection

$$\text{ParseWords}(T, T_{k-1}, T') \rightarrow \text{ParseWords}(T, T_{k-1}, T') \setminus \text{ParseWords}(T, T_{k-1}, T').$$

Consequently, $\text{ParseWords}(T, T_{k-1}, T')$ contains precisely half the parsewords of $\text{ParseWords}(T, T_{k-1})$, i.e. $3 \cdot 2^{n-1-k}$. From Theorem 7.6, one has $|\text{ParseWords}(T, T_{k-1}, T')| = |\text{ParseWords}(T, T')|$, so $|\text{ParseWords}(T, T')| = 3 \cdot 2^{n-1-k}$, as desired.

Corollary 7.8. *Since $k \leq n-2$ by Proposition 3.2, any pair of trees comparable in $\mathcal{C}_n$ has a common parse word.*

Theorem 7.9. *Suppose T_1 and T_2 have an upper bound in $\mathcal{C}_n$. Then,*

$$\text{ParseWords}(T_1, T_2) = \text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2).$$

Proof. The statement is clear when T_1 and T_2 are comparable, so assume T_1 and T_2 are not comparable. Without loss of generality, we may assume $T_1 \wedge T_2 = \text{RCT}(n)$ by repeatedly applying Proposition 7.3. Then all parenthesis pairs in RP_{T_1} and RP_{T_2} are disjoint. By Theorem 7.6,

$$\text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2, T_1) = \text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_1) = \text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2, T_2).$$

One then immediately has that $\text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2) \subset \text{ParseWords}(T_1, T_2)$.

All that remains is to show inclusion the other way. Suppose the theorem holds for trees with $k < n$ leaves. There are two cases:

- (1) Suppose T_1 and T_2 share a leaf reduction at, say, i . Then $T_1 \wedge T_2$ and $T_1 \vee T_2$ must also share this leaf reduction. Then,

$$\begin{aligned} \text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2) &= f_i^<(\text{ParseWords}(\widehat{T_1 \wedge T_2}, \widehat{T_1 \vee T_2})) \\ &\sqcup f_i^>(\text{ParseWords}(\widehat{T_1 \wedge T_2}, \widehat{T_1 \vee T_2})) \\ &= f_i^<(\text{ParseWords}(\hat{T}_1, \hat{T}_2)) \sqcup f_i^>(\text{ParseWords}(\hat{T}_1, \hat{T}_2)) \\ &= \text{ParseWords}(T_1, T_2), \end{aligned}$$

as desired, with the inductive hypothesis being used in the second inequality.

- (2) Suppose, on the other hand, that no such common leaf reduction exists. Suppose $w = w_1 w_2 \cdots w_n \in \text{ParseWords}(T_1, T_2)$. Then, since $T_1 \wedge T_2$ is $\text{RCT}(n)$, either RP_{T_1} or RP_{T_2} contains a parenthesis pair enclosing a_{n-1} , else both trees would have a leaf reduction at $(n-1, n)$. Without loss of generality, assume RP_{T_1} contains a parenthesis pair enclosing a_{n-1} . Moreover, RP_{T_1} has a maximal parenthesis pair enclosing the leaves $a_j, \dots, a_{n-1}$.

Then, since all parenthesis pairs in RP_{T_1} and RP_{T_2} are disjoint, none of $a_j, \dots, a_{n-1}$ are enclosed by a parenthesis pair in RP_{T_2} . Consequently, the subtrees of T_2 and $T_1 \wedge T_2$ with leaf set $a_j, \dots, a_n$ are both isomorphic to $\text{RCT}(n - j + 1)$. Call this subtree X_1 . By the maximality of the parenthesis pair containing $a_j, \dots, a_{n-1}$, one has that T_1 and $T_1 \vee T_2$ have isomorphic subtrees whose leaf sets are $a_j, \dots, a_n$, call this subtree X_2 . Consequently, $w_j \cdots w_n$ parses the subtree containing $a_j, \dots, a_n$ in $T_1, T_2, T_1 \wedge T_2$ and $T_1 \vee T_2$. Collapse the subtrees X_1 and X_2 to obtain $T'_1, T'_2, T'_1 \wedge T'_2$ and $T'_1 \vee T'_2$. By induction, $\text{ParseWords}(T'_1, T'_2) = \text{ParseWords}(T'_1 \wedge T'_2, T'_1 \vee T'_2)$, meaning $w_1 \cdots w_{j-1} w_{X_1} = w_1 \cdots w_{j-1} w_{X_2}$ parses $T'_1 \wedge T'_2$ and $T'_1 \vee T'_2$. It is then easy to see that this implies w lies in $\text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2)$, as desired. $\square$

Remark. If T_1 and T_2 have an upper bound in $\mathcal{C}_n$, and $\text{rank}(T_1 \vee T_2) - \text{rank}(T_1 \wedge T_2) = k$, combining Theorem 7.9 and Theorem 7.7, one has

$$|\text{ParseWords}(T_1, T_2)| = |\text{ParseWords}(T_1 \wedge T_2, T_1 \vee T_2)| = 3 \cdot 2^{n-1-k}.$$

ACKNOWLEDGEMENTS

This research was carried out at the School of Mathematics, University of Minnesota - Twin Cities, under the supervision of Vic Reiner and Dennis Stanton with funding from NSF grant DMS-1001933. We would like to thank Vic Reiner and Dennis Stanton for their invaluable guidance and support, Bobbe Cooper for giving a talk introducing us to this problem, Nathan Williams for his words of advice and continual assessment of our progress, and Alan Guo for his decisive input in the proof of Theorem 3.2.

REFERENCES

- [1] B. Cooper, E. Rowland, and D. Zeilberger. Toward a language theoretic proof of the Four Color Theorem, 2010, [arXiv:1006.1324v1 \[math.CO\]](#).
- [2] P. H. Edelman. Tableaux and chains in a new partial order of S_n . *J. Combin. Theory Ser. A*, 51(2):181–204, 1989.
- [3] S. Huang and D. Tamari. Problems of associativity: A simple proof for the lattice property of systems ordered by a semi-associative law. *J. Combinatorial Theory Ser. A*, 13:7–13, 1972.
- [4] L. H. Kauffman. Map coloring and the vector cross product. *J. Combin. Theory Ser. B*, 48(2):145–154, 1990.
- [5] D. E. Knuth. *The Art of Computer Programming*, volume 3: Sorting and Searching. Addison-Wesley Publishing Company, Reading, Massachusetts, 1973.
- [6] J. M. Pallo. Enumerating, ranking and unranking binary trees. *Comput. J.*, 29(2):171–175, 1986.
- [7] J. M. Pallo. Right-arm rotation distance between binary trees. *Inform. Process. Lett.*, 87(4):173–177, 2003.
- [8] D. D. Sleator, R. E. Tarjan, and W. P. Thurston. Rotation distance, triangulations, and hyperbolic geometry. *J. Amer. Math. Soc.*, 1(3):647–681, 1988.
- [9] R. P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997.
- [10] H. S. Wilf. *generatingfunctionology*. Academic Press Inc., Boston, MA, second edition, 1994.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455, USA
E-mail address: csar@math.umn.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, NJ 08544, USA
E-mail address: rsengupt@princeton.edu

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, MA 02139, USA
E-mail address: warutsuk@mit.edu